

Your TzielD Treatment Path

What Is TzielD?

TzielD is a prescription medicine used to delay the onset of Stage 3 type 1 diabetes, which is when your body cannot make enough insulin on its own and may require insulin injections. TzielD is for adults and children 1 year of age and older who have Stage 2 type 1 diabetes. This means that they have tested positive for 2 or more type 1 diabetes-related autoantibodies, have abnormal blood sugar levels, and do not have type 2 diabetes or other forms of diabetes.

It is not known if TzielD is safe and effective in children under 1 years of age.

How Will I Receive TzielD?

- TzielD is given by a healthcare provider through a needle placed in a vein (intravenous infusion) in the arm
- TzielD is received as an infusion one-time a day, every day, for 14 days
- If you are an adult or your child is 8 years of age or older, each TzielD infusion will last at least 30 minutes. If your child is 1 to less than 8 years of age, each TzielD infusion will last at least 2 hours. Consider planning for additional time before and after your infusion for preparation and monitoring
- For the first 5 days of treatment, your healthcare provider will give medicines by mouth before starting the TzielD infusion. These medicines include pain relievers, an antihistamine, and possibly an anti-nausea medicine. These medicines may help reduce symptoms of Cytokine Release Syndrome (CRS) such as fever, headache, muscle and joint pain, or nausea, with or without vomiting. Vomiting and diarrhea may occur more frequently in children less than 8 years of age

If a scheduled infusion is missed, your healthcare provider will continue the treatment on the next scheduled day. Two infusions are not given on the same day.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about TzielD?

TzielD may cause serious side effects, including:

- **Viral Reactivation.** Epstein-Barr virus (EBV) and cytomegalovirus (CMV) are common viruses that may stay inactive in your body after an initial infection. TzielD may cause these viruses to become active again which, especially in patients with a weakened immune system, can become serious and potentially life-threatening. These viral reactivations can happen during treatment with TzielD and up to 2 months after your last dose. Your healthcare provider will test you for active EBV and CMV infections before treatment with TzielD. Contact your healthcare provider right away if you develop symptoms of an infection during or after treatment with TzielD (such as fever, swollen glands, or fatigue).

Please see additional Important Safety Information below and full [Prescribing Information](#), including Risk of **SERIOUS SIDE EFFECTS** and [Medication Guide](#).


Your TzielD Treatment Path

You and your doctor have decided that TzielD is the right choice. It's now time to take the next steps. Whether you're a patient or a caregiver, this guide will walk you through what to expect, from selecting an infusion site and enrolling in TzielD COMPASS—the TzielD patient support program—to understanding insurance coverage and scheduling your infusions.

TzielD COMPASS is here to support you every step of the way, helping to make your treatment path as smooth as possible.

Let's get started.


Starting TzielD and Insurance Approval



Consider a Site of Care (SOC) and Infusion Method

- Identify infusion locations (office, home, or both) that work with your insurance coverage, location, transportation, and scheduling needs
- Talk to your doctor about different infusion methods, such as a daily IV infusion, a midline, or a peripherally inserted central catheter (PICC)
- Enroll and share your choices with TzielD COMPASS, which will work with your health plan to review coverage and help your doctor support the approval process

Your doctor will explain your treatment options and help you decide which method and location are best for you.



Enroll in TzielD COMPASS

- Learn about TzielD COMPASS and the ways the program can help throughout your or your loved one's treatment journey
- Provide your signature or eConsent to complete your TzielD COMPASS enrollment
- Connect with a Clinical Educator (CE) to learn more about what to expect during your treatment. A CE can also connect you with a TzielD Mentor, a real patient or caregiver who has experienced TzielD treatment firsthand

Your doctor will complete the enrollment form with your help, including asking for needed information and your final consent to enroll.

Who Can Help?

COMPASS Navigators are available throughout your treatment journey **whenever you have questions** or need additional resources or support.

You can reach out to your **CE** if you have questions about the infusion process.




Understand Your Insurance Coverage

- Review your insurance coverage and benefits with your COMPASS Navigator
- Make a final decision with your doctor about your SOC and infusion options, based on your insurance coverage
- Learn what financial assistance may be available

Based on your coverage options, your doctor can help you make your final infusion decisions.

TzielD Treatment and Follow-Up



Schedule Your Treatment

- Work with your doctor to schedule your 14-day treatment
- Your doctor will provide instructions on when to take your pre-infusion medications and complete any pre-infusion lab work
- Ask your doctor any remaining questions you may have about your infusion

Your doctor will help you schedule your infusion and finish preparing for treatment.



Complete Your Treatment

- Confirm your infusion appointment through your COMPASS Navigator
- Complete all 14 daily infusions and receive check-ins from your doctor and TzielD COMPASS Navigator along the way
- A CE is available to help coordinate a second check-in with your TzielD Mentor during your infusion

During your treatment period, your doctor will check in with you to make sure everything is going okay.



Begin Post-Treatment Follow-Up

- Discuss post-treatment monitoring and care with your doctor
- Contact your COMPASS Navigator if you have any insurance or coverage questions

Once treatment is done, your doctor will meet with you and review your follow-up care.

Learn more

Click [here](#) to see how you can **receive personalized support** along your treatment path.

If you have any questions, please call TzielD COMPASS at 1-844-778-2246,

Monday through Friday from 8:00 AM–8:00 PM ET, or send us an email at COMPASS@sanofi.com

IMPORTANT SAFETY INFORMATION (cont'd)

What is the most important information I should know about TzielD? (cont'd)

TzielD may cause serious side effects. These include (cont'd):

- **Cytokine release syndrome (CRS).** Signs and symptoms may start during the first 5 days of TzielD treatment and could include fever, nausea (with or without vomiting), feeling tired (fatigue), headache, muscle and joint pain, or increased liver enzymes in your blood. Tell your healthcare provider right away if you develop any signs and symptoms of CRS during treatment with TzielD.
- **Decrease in white blood cells.** TzielD may cause a decrease in a type of white blood cell called lymphocytes. A decrease in white blood cells is a serious, but common side effect that can affect your body's ability to fight infections. A decrease in white blood cell counts can happen after your first dose. Your white blood cell counts will start to go back to normal after your fifth dose of TzielD. Some people may develop longer and more severe decreases in lymphocytes.

Your healthcare provider will do blood tests to check for active infections, verify your liver function and your complete blood counts before you start treatment and during treatment with TzielD. During and after your treatment with TzielD, your healthcare provider will check for serious side effects, as well as other side effects, and treat you as needed. Your healthcare provider may temporarily or completely stop your treatment with TzielD, if you develop liver problems, have a serious infection or viral reactivation, or if your blood counts stay too low.

What should I tell my healthcare provider before receiving TzielD?

Before or after receiving TzielD, tell your healthcare provider about all your medical conditions, including if you:

- have any of the conditions or symptoms listed in the section "What is the most important information should know about TzielD?"
- have a weakened immune system, including if you have Down syndrome.

- have a serious infection or an infection that does not go away or keeps coming back (chronic).

- are immunizing (vaccine). TzielD may affect how well a vaccine works. Tell your healthcare provider that you are receiving treatment with TzielD before receiving a vaccine.

- are pregnant or plan to become pregnant. TzielD may harm your unborn baby. **Do not** receive TzielD during pregnancy and at least 30 days before a planned pregnancy.

If you become pregnant while taking TzielD, you are encouraged to report your pregnancy to us at our Adverse Event reporting line at 1-800-633-1610 or visit <https://ae.reporting.sanofi>.

- are breastfeeding or plan to breastfeed. It is not known if TzielD passes into your breast milk and if it can harm your baby. Talk to your healthcare provider about the best way to feed your baby if you receive TzielD.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of TzielD?

The most common side effects of TzielD include:

- decrease in white blood cell counts
- vomiting
- rash
- diarrhea
- headache

These are not all of the possible side effects of TzielD. For more information, ask your healthcare provider or pharmacist. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088. You may also report side effects to us at 1-800-633-1610 or visit <https://ae.reporting.sanofi>.

Please see full [Prescribing Information](#), including Risk of **SERIOUS SIDE EFFECTS** and [Medication Guide](#).